

Running title: Chemical Composition and Microbial Activities of Dried Bush Tea

Effect of Different Drying Methods on Chemical Composition and Microbial Activities of Bush Tea (*Athrixia phylicoides*)

Fhatuwani N. Mudau* and Wonder Ngezimana

Department of Agriculture and Animal Health, College of Agriculture and Environmental Sciences, University of South Africa, Private Bag X6, Florida, 1710, South Africa

*For correspondence: mudaufn@unisa.ac.za

Novelty statement

There is need for research in best bush tea drying methods to enable commercialization of this plant with beneficial phytochemicals, antioxidants and microbial activities. The current research was attributed by the fact that there is no drying protocol which has already been published on bush tea and the current work will pave ways on future herbal bush tea industry.

Abstract

The effect of different drying methods; sun, freeze, shade and oven drying on total polyphenol content, tannins, total antioxidants and phytochemicals content of bush tea (*Athrixia phylicoides* DC.) were evaluated. Assessment of minimum inhibitory concentration (MIC) assay was also done on bush tea samples. Results showed that different drying processes significantly affected phytochemical compositions of bush tea. Highest total phenolic content (8.34 mg/100 g) sample was found on freeze and shade dried bush tea compared to less than (6.50 mg/100 g) in sun or oven dried samples. Similar response was shown in total antioxidants with shade (0.628 µmol/g) and freeze (0.626 µmol/g) drying having significantly highest content of total antioxidants as compared to sun (0.440 µmol/g) and oven (0.444 µmol/g) dried samples. MIC values falls within 3.1 to 6.3 mg/mL with gram-positive bacteria being more vulnerable to the inhibitory effect of extract regardless of bush tea drying

method than gram-negative. Anti-microbial activities of bush tea observed in this study will be useful to develop tea extracts for deterrence of diseases causing microorganisms hence shade drying methods will provide easy drying bush tea samples which can be adopted by farmers for commercial purposes.

Keywords: Antioxidants; Bush tea; Drying methods; Minimum inhibitory concentration; Phytochemicals

Introduction

Bush tea (*Athrixia phylicoides* DC.) leaves contains 5-hydroxy-6,7,8,3',4',5'-hexamethoxy flavon-3-ol (Mashimbye *et al.*, 2006), 3-O-demethyldigicitrin, 5,6,7,8,3',4'-hexamethoxyflavone and quecetin (Mavundza *et al.*, 2010), tannins (Mudau *et al.*, 2007; Chabeli *et al.*, 2008), total polyphenols (Mudau *et al.*, 2006; Maudu *et al.*, 2012) and total antioxidants (Maudu *et al.*, 2010; Mogotlane *et al.*, 2007) compounds with antimicrobial activity (Mavundza *et al.*, 2010). Indigenous South Africans have significant use of this plant as a traditional medicine (Roberts, 1990; Van Wyk and Gericke, 2000; Mbambezeli, 2005).

McGaw *et al.* (2007) reported that bush tea contains no caffeine as well as pyridoxine making it more appropriate as a healthy beverage. More so, occurrence of antioxidants in bush tea has positive favourable health benefits (Mudau and Mariga, 2012). According to Mudau *et al.* (2006), active chemical compounds present in herbal tea influence the quality of herbal tea. They also serve as the potential indicators of medicinal prospective due to their antioxidant activities (Mudau *et al.*, 2006; MaKay and Blumberg, 2007; Kokotkiewicz and Luczkiewicz, 2009). However, yield and chemical composition from herbal plants are related to a variety of factors which include the drying process (Rocha *et al.*, 2011).

Drying is an important part of tea post-harvest handling, with the processing having impact on antioxidant content and appearance which all have effect on viable of the industry (Chong and Lim, 2012). It is the most common and fundamental way to preserve quality of aromatic and medicinal plants (Müller and Heindl, 2006; Rocha *et al.*, 2011). In other common herbal tea in South Africa, Joubert and de Villiers (1997) reported that drying affects quality attributes on rooibos tea whilst in

honeybush tea, drying methods did not significantly influence the quality of tea (du Toit and Joubert, 1998). In other herbal teas, sun-drying resulted in deterioration of antioxidant properties (Chong and Lim, 2012) and various drying methods have different effects on aromatic and medicinal plants (Stafford *et al.*, 2005; Rocha *et al.*, 2011). Hence for viability of the bush tea production, there is need to establish drying techniques for recommendations to producers.

Data that describe the standard production protocol for drying bush tea have not been established. Therefore, the objective of this study was to compare effect of different bush tea drying methods on phytochemical composition and anti-microbial activities of bush tea extract.

Materials and Methods

Fresh bush tea samples were collected from the wild at Muhuyu village (22°53'60S and 30°25'0E, 724 m.a.s.l.) in Limpopo province, South Africa. The fresh leaves of uniform shape, color and size were selected and subjected to four different drying methods viz., sun, freeze, shade and oven drying. In sun drying, bush tea leaves were placed in the sun/daylight. Midday temperature reached 35°C for 5 days during the sun drying period. For, freeze drying, leaves were placed in freezer dryer. In shade drying, leaves were placed in wooden trays and protected from direct sunlight. For oven drying, leaves were placed in an oven tray and heated at a temperature between 45 to 65°C overnight. All treatment were laid out in a completely randomised design (CRD) and replicated five times. Dry samples were thereafter analysed for chemical compositions. Assessment of minimum inhibitory concentration (MIC) assay was also done (Eloff, 1988). Comparisons of total phenolic contents and antioxidant activities were made to the fresh bush tea extracts reported by Mavundza (2010).

Chemical Compositions Assays

Determinations of total polyphenol content were done using methods reported by Singleton and Rossi (1965) and modified by Waterman and Mole (1994). Tannins were determined using Vanillin HCL method described by Prince *et al.* (1978) and total antioxidants were determined using the method described by Awika *et al.* (2004).

Phytochemicals Assays

The chlorophyll, riboflavin, niacin, ascorbic acid, carotenoid contents were determined using the standard AOAC methods (1984).

Minimum Inhibitory Concentration Assay

Gram-positive; *Staphylococcus aureus* (ATCC 12600), *Bacillus cereus* (ATCC 11778), *B. subtilis*, *B. pumilis* (ATCC 21356), *Enterococcus faecalis* (ATCC 29212) and Gram-negative; *Pseudomonas aeruginosa* (ATCC 25922), *Escherichia coli* (ATCC 11775), *Klebsiella pneumonia* (ATCC 27736) microorganisms were used. The assays were carried out using micro-dilution methods on 96 well microplates (Eloff, 1988).

Data Analysis

Data were subjected to analysis of variance (ANOVA) using PROC GLM (General linear model) procedure of SAS version 9.2 (SAS Int, 2011).

Results

Chemical Compositions

Drying of bush tea after harvesting had significant effect on the chemical composition of the extract, with generally a higher composition than in fresh samples which had 2.30 mg/100 g of total phenolic content. Total phenolics were higher in shade (8.34

mg/100 g) and freeze (8.34 mg/100 g) dried bush tea as compared to sun (6.42 mg/100 g) and oven (5.62 mg/100 g) dried samples (Table 1). The same response was shown in total antioxidants with shade (0.63 $\mu\text{mol/g}$) and freeze (0.63 $\mu\text{mol/g}$) drying having significantly highest content of total antioxidants as compared to sun (0.44 $\mu\text{mol/g}$) and oven (0.44 $\mu\text{mol/g}$) dried samples. However, even though there was slightly more tannin content (0.4 mg/100 g) in freeze dried bush tea samples than in other drying techniques; the different drying approaches did not exhibit significant differences on tannin contents of bush tea leaves.

Shade (83.4 mg/100 g) and freeze (83.4 mg/100 g) dried bush tea samples had significantly highest chlorophyll contents compared to sun (69.9 mg/100 g) and oven (68.3 mg/100 g) dried samples (Table 2). The difference between the highest and lowest contents was 14.1 mg/100 g of dried sample. The ascorbic acid, niacin, carotenoids and riboflavin contents of the sampled bush tea did not vary regardless of the drying method used. Average phytochemical compositions within the plant samples were 76.3 mg/100 g, 1.7 $\mu\text{mol/g}$, 1.5 mg/100 g, 1.8 mg/100 g and 2.4 mg/100 g for chlorophyll, ascorbic acid, niacin, carotenoids and riboflavin contents respectively.

Minimum Inhibitory Concentration

MIC values of extract on different microorganisms are shown in Table 3. The crude extract showed positive inhibitory activity against the entire gram-positive and gram-negative tested microorganism regardless of drying method subjected to the bush tea sample with MIC values ranging from 3.1 to 6.3 mg/mL. All the gram-negative microorganisms gave a MIC value of 6.3 mg/mL whilst the lowest MIC values of 3.1 mg/mL were obtained in gram-positive microorganisms except for *B. cereus* with MIC value of 6.3 mg/mL.

Discussion

Drying of bush tea resulted in a higher composition of chemical constituents. During drying, cell membranes rupture and degrade thereby releasing compounds during extraction (Stafford *et al.*, 2005). However, freeze and shade drying proved to be more efficient in retaining the total polyphenols and antioxidants within the bush tea samples than oven and sun drying procedures. In most studies carried out on various aromatic and medicinal plant types (Yousif *et al.*, 2000; Asami *et al.*, 2003; Abascal *et al.*, 2005), freeze drying has been shown as mostly recommended method in retention of plant compounds. Similarly, van Golde *et al.* (2004) reported that approximately 70% of polyphenols in red wine can be preserved through freeze drying. In grape peel, losses of polyphenols, tannins and antioxidants have been reported in oven dried samples at higher temperature (Laurrauri *et al.*, 1997).

Post-harvest processes like drying take authoritative impact on the quality of the product influencing its value (Müller and Heindl, 2006) which is mostly shown on its appearance. The present study showed that sun and oven drying can reduce the chlorophyll content of bush tea, adversely affecting the color. Mahanom *et al.* (1999) observed loss of chlorophyll in all oven dried medicinal plants leaves. Similar results were reported by Joubert and de Villiers (1997) in rooibos tea, where less chlorophyll was present in sun-dried than in tea dried under controlled conditions. Ramana *et al.* (1988) suggested that such loss of colour must be minimized as it may adversely distance prospective customers due to possibility of below par processing. In addition to yielding extracts high in phenols and active ingredients; meadowsweet (*Filipendula ulmaria*) and willow (*Salix alba*) herbs developed an appropriate colour

for fusion into a beverage when dried under temperatures of 30°C compared to the 70°C (Harbourne *et al.*, 2009).

Besides the benefits of antioxidants, bush tea samples were also beneficial as antibacterial regardless of drying method used in this trial. According to Fabry *et al.* (1998), all plant extracts with MIC values below 8 mg/mL are considered to possess some antimicrobial activity. MIC values have demonstrated that the gram-positive bacteria appeared to be more susceptible than the gram-negative ones to the inhibitory effect of the extract. Similar results were reported by Tshikalange *et al.* (2005) who also noted on different susceptibility of these bacteria while studying the antibacterial activities of other herbal plants. These difference in susceptibility can be ascribed to the morphological differences between the gram-positive and gram-negative bacteria (Palombo and Semple, 2001; Tadeg *et al.*, 2005), with the later having an outer phospholipidic membrane that carries structural lipopolysaccharide components making the cell walls impermeable to lipophilic solutes (Nostro *et al.*, 2000).

Benefits of tea extracts have been noted previously by Toda *et al.* (1989), with the extract inhibiting enteric pathogens such as *Staphylococcus aureus*, *S. epidermis* and *Plassiomonas shigelloides*. Amongst other things, tea extracts have been found effective against *Helicobacter pylori* which are linked to gastric, peptic and duodenal ulcer diseases (Diker and Hascelik, 1994). Biological activities of the cariogenic streptococci can also be inhibited by polyphenols in green tea thereby preventing teeth from decaying (Sakanaka *et al.*, 1996; Mitscher *et al.*, 1997). According to Baydar *et al.* (2004), the extent of this inhibition can be attributed to phenolic composition within the plant with phenols as the predominant active compounds in

medicinal plants (Rios and Recio, 2004). This makes bush tea a very promising herbal tea as it is a rich source of phenols.

In conclusion, shade and freeze drying methods have shown to be more useful for retention of phytochemicals within bush tea samples. Crude extracts of bush tea had a broad spectrum in their actions on antibacterial activities of either gram positive bacteria or gram negative bacteria regardless of drying method subjected to the bush tea. Therefore, sound processed bush tea could be used as accessible sources of natural antimicrobial and antioxidants.

References

- Abascal, K., L. Ganora and E. Yarnell, 2005. The effect of freeze-drying and its implications for botanical medicine: A review. *Phytother. Res.*, 19: 655–660
- AOAC, 1984. Vitamins and other nutrients. In *Official Methods of Analysis of the Association of Official Analytical Chemists*, pp 838 – 841. 14th Edition (William S, ed.), AOAC, Virginia
- Asami, D.K., Y.J. Hong, D.M. Barrett and A.E. Mitchell, 2003. Comparison of the total phenolic and ascorbic acid content of freeze dried and air-dried marionberry, strawberry, and corn grown using conventional, organic and sustainable agricultural practices. *J. Agric. Food Chem.*, 51: 1237–1241
- Awika, J.M., L.W. Rooney and R.D. Waniska, 2004. Anthocyanins from black sorghum and their antioxidant properties. *J. Food Chem.*, 90: 293–301
- Baydar, N. G., G. Ozkan, and O. Sagdiç, 2004. Total phenolic contents and antibacterial activities of grape (*Vitis vinifera* L.) extracts. *Food Control.*, 15: 335–339

- Chabeli, P.M., F.N. Mudau, P.W. Mashela and P. Soundy, 2008. Response of tannin content of bush tea (*Athrixia phylicoides* DC.) to nitrogen, phosphorus and potassium nutrition as affected by season. *S. A. J. Plant Soil.*, 25: 79–83
- Chong, K.L. and Y.Y. Lim, 2012. Effects of drying on the antioxidant properties of herbal tea from selected *Vitex* species. *J. Food Quality.*, 35: 51–59
- Diker, K.S. and G. Hascelik, 1994. The bactericidal activity of tea against *Helicobacter pylori*. *Lett. App. Microbiol.*, 19: 299–300
- du Toit, J. and E. Joubert, 1998. Effect of drying conditions on the quality of honeybush tea (*Cyclopia*). *J. Food Processing Preservation.*, 22: 493–507
- Eloff, J.N., 1998. A sensitive and quick microplate method to determine the minimum inhibition concentration of plant extracts for bacteria. *Planta Medica.*, 64: 711–713
- Fabry, W., Okemo, P. O., and Ansorg, R. 1998. Antibacterial activity of East African medicinal plants. *J. Ethnopharmacol.*, 60: 79–84
- Harbourne, N., E. Marete, J.C. Jacquier and D. O’riordan, 2009. Effect of drying methods on the phenolic constituents of meadowsweet (*Filipendula ulmaria*) and willow (*Salix alba*). *Food Sci. Technol.*, 42: 1468–1473
- Joubert, E. and O. T. de Villiers, 1997. Effect of fermentation and drying conditions on the quality of rooibos tea. *Inter. J. Food Sci. Technol.*, 32: 127–134
- Kokotkiewicz, A. and M. Luczkiewicz, 2009. Honeybush (*Cyclopia* sp.) - a rich source of compounds with high antimutagenic properties. *Fitoterapia.*, 80:3–11

- Laurrauri, J.A., P. Ruperez and F. Saura-Calixto, 1997. Effect of drying temperature on the stability of polyphenols and antioxidant activity of red grape pomace peel. *J. Agric Food Chem.*, 45: 1390–1393
- Mahanom, H., A.H. Azizah and M.H. Dzulkifly, 1999. Effect of different drying methods on concentrations of several phytochemicals in herbal preparation of 8 medicinal plants leaves. *Mal. J. Nutr.*, 5: 47–54
- MaKay, D.L. and J.B. Blumberg, 2007. A review of the bioactivity of South African herbal teas: rooibos (*Aspalathus linearis*) and honeybush (*Cyclopia intermedia*). *Phytother. Res.*, 21: 1–16
- Mashimbye, M.J., F.N. Mudau, P. Soundy and T. Van Ree, 2006. A new flavonol from *Athrixia phyllicoides* (Bush tea). *S. A. J. Chem.*, 59: 1–2
- Maudu, M., F.N. Mudau and I.K. Mariga, 2010. The effect of pruning on growth and chemical composition of cultivated bush tea (*Athrixia phyllicoides* D.C). *J. Med. Plants Res.*, 4: 2353–2358
- Maudu, M., F.N. Mudau and I.K. Mariga, 2012. Quality profiles of cultivated and wild bush tea (*Athrixia phyllicoides*) harvested at various phenological stages. *Int. J. Agric. Biol.*, 14: 552–556
- Mavundza, E.J., T.E. Tshikalange, N. Lall, A.A. Hussein, F.N. Mudau and J.J. Meyer, 2010. Antioxidant activity and cytotoxicity effect of flavonoids isolated from *Athrixia phyllicoides*. *J. Med. Plants Res.*, 4: 2584–2587
- Mavundza, E.J., 2010. Antioxidant and antibacterial activities of ethanol extract and flavonoids isolated from *Athrixia phyllicoides*. MSc Thesis. University of Pretoria. South Africa

- Mbambezeli, G., 2005. *Athrixia phylicoides*. South African National Biodiversity Institute
- McGaw, L.J., V. Steenkamp and J.N. Eloff, 2007. Evaluation of *Athrixia* bush tea for cytotoxicity, antioxidant activity, caffeine content and presence of pyrrolizidine alkaloids. *J. Ethnopharmacol.*, 110: 16–22
- Mitscher, L.A., M. Jung, D. Shankel, J.H. Dou, L. Steele and S. Pillai. 1997. Chemoprotection: a review of the potential therapeutic antioxidant properties of green tea (*Camellia sinensis*) and certain of its constituents. *Med. Res. Rev.*, 17: 327–365
- Mogotlane, I.D., F.N. Mudau, P.W. Mashela and P. Soundy, 2007. Seasonal responses of total antioxidant contents in cultivated bush tea (*Athrixia phylicoides* L.) leaves to fertilizer rates. *Med. Aromatic Plant Sci. Biotech.*, 1: 77–79
- Mudau, F.N. and I.K. Mariga, 2012. Bush tea as a herbal beverage and medicinal plant. In V.R. Reedy, (ed), *Tea in Health and Disease Prevention.*, 1st Edition, pp 182–192. Elsevier. Academic press, Oxford. UK
- Mudau, F.N., H.T. Araya, E.S. du Toit, P. Soundy and J. Olivier, 2007. Bush Tea (*Athrixia phylicoides* DC.) as an alternative herbal and medicinal plant in Southern Africa: Opportunity for commercialization. *Med. Aromatic Plant Sci. Biotech.*, 1: 70–73
- Mudau, F.N., P. Soundy, E.S. Du Toit and J. Oliver, 2006. Variation in polyphenolic content of *Athrixia phylicoides* (L.) (bush tea) leaves with season and nitrogen application. *S. A. J. Bot.*, 72: 398–402

- Müller, J. and A. Heindl, 2006. Drying of medicinal plants. In R.J. Bogers, L.E. Craker and D. Lange (eds.), *Medicinal and Aromatic Plants*, pp 237–252. Springer. Netherlands
- Nostro, A., M.V. Germanó, V.D. Angelo, A. Marino and M.A. Cannatelli, 2000. Extraction methods and bioautography for evaluation of medicinal plant antimicrobial activity. *Lett. App. Microbiol.*, 30:379–384
- Palombo, E. A. and S. J. Semple, 2001. Antibacterial activity of traditional Australian medicinal plants. *J. Ethnopharmacol.*, 77: 151–157
- Prince, M.L., S. Van Scoyoc and L.G. Butler, 1978. Critical evaluation of the vanillin reaction as an assay for tannin in sorghum grain. *J. Agric. Food Chem.*, 26: 1214–1218
- Ramana, S.V., K.S. Jayaraman and B.L. Mohan Kumar, 1988. Studies on the colour of some dehydrated green leafy vegetables. *Indian Food Packer.*, ¾: 19–23
- Rios, J.L. and M.C. Recio, 2005. Medicinal plants and antimicrobial activity. *J. Ethnopharmacol.*, 100: 80–84
- Roberts, M., 1990. *Indigenous healing plants*, pp. 51–52. Southern Book publishers: Half way House, South Africa
- Rocha, R.P., E.C. Melo and L.L. Radünz, 2011. Influence of drying process on the quality of medicinal plants: A review. *J. Med. Plants Res.*, 5: 7076–7084
- Sakanaka, S., M. Aizawa, M. Kim and T. Yamamoto, 1996. Inhibitory effects of green tea polyphenols on growth and cellular adherence of an oral bacterium, *Porphyromonas gingivalis*. *Biosci. Biotechnol. Biochem.*, 60: 745–749

- Singleton, V.L. and J.A. Rossi, 1965. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American J. Enol. Viticul.*, 16: 144–158
- Stafford, G.I., A.K. Jäger and J. Van Staden, 2005. Effect of storage on the chemical composition and biological activity of several popular South African medicinal plants. *J. Ethnopharmacol.*, 97: 107–115
- Statistical Analysis System (SAS) Institute Inc., 2011. *User's Guide, Version 9.2*, Cary NC, USA
- Tadeg, H., E. Mohammed, K. Asres and T.G. Mariam. 2005. Antimicrobial activities of some selected traditional Ethiopian medicinal plants used in the treatment of skin disorder. *J. Ethnopharmacol.*, 100: 168–175
- Toda, M., S. Okubo, R. Hiyoshi and S. Tadakatsu, 1989. The bactericidal activity of tea and coffee. *Lett. App. Microbiol.*, 8: 123–125
- Tshikalange, T.E., J.J. Meyer and A.A. Hussein, 2005. Antimicrobial activity, toxicity and the isolation of a bioactive compound from plants used to treat sexually transmitted diseases. *J. Ethnopharmacol.*, 96: 515–519
- van Golde, P.H., M. van der Westelaken, B.N. Bouma and A. van de Wiel, 2004. Characteristics of piraltin, a polyphenol concentrate, produced by freeze-drying of red wine. *Life Sci.*, 74: 1159–1166
- Van Wyk, B.E. and M. Gericke, 2000. *People's Plants. A guide to useful plants of Southern Africa*, pp. 102–103. Briza Publications, Pretoria, South Africa

Waterman, P.G. and S. Mole, 1994. *Analysis of Plant Metabolites*, pp: 1–103.
Oxford: Alden Press Limited

Yousif, A.N., T.D. Durance, C.H. Scaman and B. Girard, 2000. Headspace volatiles and physical characteristics of vacuum-microwave, air and freeze-dried oregano (*Lippia berlandieri* Schauer). *J. Food Sci.*, 65: 926–930

Table1: Response of chemical composition to selected drying methods of bush tea

Drying methods	Total polyphenols content (mg/100g)	Total antioxidants ($\mu\text{mol/g}$)	Tannin contents (mg/100g)
Sun drying	6.42 b	0.44 b	0.34 a
Shade drying	8.34 a	0.63 a	0.34 a
Oven drying	5.62 b	0.44 b	0.34 a
Freeze drying	8.34 a	0.63 a	0.38 a
Significant level	0.0001	0.0001	0.5038
LSD 5%	1.01	0.08	0.07

Means in a column followed by the same letter are not significantly different ($P>0.05$).

Table 2: Effect of drying on phytochemical compositions of bush tea

Drying methods	Chlorophyll (mg/100g)	Ascorbic acid ($\mu\text{mol/g}$)	Niacin (mg/100g)	Carotenoids (mg/100g)	Riboflavin (mg/100g)
Sun drying	69.9 b	1.8 a	1.6 a	1.8 a	1.9 a
Shade drying	83.4 a	1.8 a	1.7 a	1.9 a	2.6 a
Oven drying	68.3 b	1.6 a	1.4 a	1.8 a	2.6 a
Freeze drying	83.4 a	1.6 a	1.4 a	1.8 a	2.4 a
Significant level	0.0001	0.9794	0.8969	0.9981	0.3195
LSD 5%	4.2	1.6	1.0	1.1	0.9

Means in a column followed by the same letter are not significantly different ($P>0.05$).

Table 3: MIC values (mg/ml) of the crude extract from bush tea dried irrespective of drying methods

Bacterial species	Gram +/-	Minimum inhibitory concentration (mg/ml)
<i>Staphylococcus aureus</i>	+	3.1
<i>Bacillus cereus</i>	+	6.3
<i>Bacillus subtilis</i>	+	3.1
<i>Bacillus pumilis</i>	+	3.1
<i>Enterococcus faecalis</i>	+	3.1
<i>Pseudomonas aeruginosa</i>	-	6.3
<i>Escherichia coli</i>	-	6.3
<i>Klebsiella pneumonia</i>	-	6.3