



Magnitude of non-adherence to antiretroviral therapy and associated factors among adult people living with HIV/AIDS in Benishangul-Gumuz Regional State, Ethiopia

Fikadu Tadesse Nigusso^{1,2} and Azwihangwisi Helen Mavhandu-Mudzusi¹

¹ Department of Health Studies, University of South Africa (UNISA), Pretoria, South Africa

² Nutrition and Education Section, United Nations World Food Programme (WFP), Addis Ababa, Ethiopia

ABSTRACT

Introduction. Following global efforts to increase antiretroviral therapy (ART) access and coverage, Ethiopia has made significant achievement with a 6.3% annual decline in the HIV/AIDS incidence rate between 1990 and 2016. Such success depends not only on access to ART but also on attaining optimum treatment adherence. Emerging studies in Ethiopia has shown the increasing prevalence of poor adherence and lack of the desired viral suppression, but the extent and factors associated with non-adherence to ART are not well known, especially in the current study setup. In this study, we examined the magnitude and factors associated with treatment and non-adherence to ART among people living with HIV in Benishangul-Gumuz Regional State, northwest Ethiopia.

Methods. An institutional facility based cross-sectional descriptive study was carried out among adult people living with HIV/AIDS from mid-December 2016 to February 2017 with only 98.9% response rate. Sociodemographic factors (age, gender, marital status and residential area), economic factors (educational status, income, asset possession, employment status, dietary diversity, nutritional status and food security), and clinical characteristics (CD4 count, duration on ART and history of opportunistic infections) were explanatory variables. ART non-adherence was measured using a visual analogue scale (VAS). We used binary logistic regression and subsequent multivariate logistic regression analysis to determine the factors associated with ART non-adherence.

Result. Overall, 39.7% of the participants were found non-adherent to ART. Strong association was found between non-adherence to ART and young age below 25 years (AOR: 4.30, 95% CI [1.39–3.35]; $p = 0.011$), urban residential area (AOR: 2.78, CI [1.23–7.09], $p = 0.043$), lack of employment (AOR: 1.75, 95% CI [1.05–2.91], $p = 0.032$), food insecurity (AOR: 2.67, 95% CI [7.59–8.97]; $p < 0.0001$), malnutrition (AOR: 1.55, 95% CI [1.94–2.56]; $p = 0.027$) and opportunistic infections (AOR: 1.81, 95% CI [1.11–2.97]; $p = 0.018$).

Conclusion. The prevalence of non-adherence to ART in this study was high. Sociodemographic and economic factors such as young age of below 25 years, urban residential area, lack of employment, food insecurity, malnutrition and opportunistic infections were among the factors associated with non-adherence to ART.

Submitted 12 August 2019

Accepted 13 January 2020

Published 11 August 2020

Corresponding author
Fikadu Tadesse Nigusso,
fike1f@gmail.com

Academic editor
Morenike Folayan

Additional Information and
Declarations can be found on
page 14

DOI 10.7717/peerj.8558

© Copyright
2020 Nigusso and Mavhandu-
Mudzusi

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Epidemiology, Global Health, HIV, Infectious Diseases, Public Health

Keywords Adherence, Antiretroviral therapy, HIV/AIDS, Visual analogue scale

INTRODUCTION

Antiretroviral therapy (ART) has played a significant role in responding to HIV/AIDS epidemics (*Granich et al., 2012; Williams, Lima & Gouws, 2011*). It has increased survival and improved quality of life, and reduced the rate of disease progression and death (*Oguntibeju, 2012; Dalhatu et al., 2016*). Consequently, ART service expansion has received significant support at the global and national level through the commitment of the Sustainable Development Goals (SDG) (*UNAIDS, 2015; The Federal Democratic Republic of Ethiopia Ministry of Health, 2015*). The SDG response sets out targets for HIV treatment: 90% of people living with HIV know their HIV status, 90% of people who know their status receiving treatment and 90% of people on treatment having a suppressed viral load (*UNAIDS, 2011*). In line with this initiative, HIV treatment and care services have expanded dramatically (*UNAIDS, 2018*). In 2017, over 21.7 million people living with HIV were receiving ART globally (*UNAIDS, 2018*). Regardless of this increase, ensuring adherence to HIV treatment remains challenging for the HIV/AIDS response, mostly in sub-Saharan Africa (*Fonsah et al., 2017; Eyassu, Mothiba & Mbambo-Kekana, 2016; Wang et al., 2011*) including Ethiopia (*Alagaw et al., 2013; Peltzer & Pengpid, 2013*).

Scientific literature indicates that for ART to be effective and prevent the emergence of resistant strains, a strict adherence level of $\geq 95\%$ is recommended. This was based on Paterson's pioneer study that found up to 95% adherence is necessary for effective HIV viral suppression (*Paterson et al., 2000*). Once initiated, ART must be continued daily (*World Health Organization, 2010*). However, irrespective of the benefits of good adherence, not all people living with HIV/AIDS (PLWHA) fully adhere to the required treatment level. Failure to attain the required adherence level results in poorer prognosis, higher morbidity, mortality and the development of resistance to ART (*Iacob, Iacob & Jugulete, 2017; Nachege et al., 2011*). Once resistance develops, it has dire consequences for individual PLWHA, family, community, healthcare providers and the healthcare system because treating ART resistant HIV strain requires second-line ART regimens, that are more expensive and have worse side effects (*Chauhan et al., 2019; Karade et al., 2018*).

Given the importance of adherence, a growing body of research has identified factors associated with non-adherence to ART as behaviors related to the patient itself, therapy related (medication regimens such as more than one tablet or dosing time per day and ART toxicity), poor relationships with healthcare providers and socioeconomic factors (*Kim et al., 2018; Mohammed, Ahmed & Tefera, 2015; Wasti et al., 2012*). Poor ART adherence was reported among underprivileged communities where socioeconomic inequalities heighten the risk (*Young et al., 2014*) and lower quality of life is prevalent (*Liping et al., 2015; Hansana et al., 2013*), negatively impacting the HIV program (*Wang et al., 2011*). For PLWHA, factors related to gender dimensions, socioeconomic factors such as education status, income, asset possession and food security have been associated with non-adherence to ART (*Basti et al., 2017; Abera et al., 2015; Alagaw et al., 2013*). For example, in South

Africa (*Eyassu, Mothiba & Mbambo-Kekana, 2016*), Cameroon (*Fonsah et al., 2017*) and northern Tanzania (*Samuel Edward et al., 2018*), gender, income and level of education were reported as the determinants of ART adherence. Another study in Wolaita Sodo, Ethiopia, found lack of food was associated with poor adherence (*Alagaw et al., 2013*). In opposition to the culminated evidence, there are studies that have reported an absence of the impacts of socioeconomic inequalities on ART adherence in middle and low-income countries (*Peltzer & Pengpid, 2013*).

Ethiopia is among sub-Saharan African countries sturdily hit by epidemics. The country has made a significant achievement in responding to the HIV/AIDS epidemic, with a 6.3% annual decline in the HIV/AIDS incidence rate between 1990 and 2016 and a total reduction of 77% (*Deribew et al., 2019*). Regardless of the government's commitment to end the HIV/AIDS epidemic as per the 2030 SDG initiative (particularly one that envisages a 90% suppression of viral loads to prevent treatment failure and reduce AIDS-related deaths), achieving the required level of suppressed viral load is becoming a challenge in Ethiopia. Current evidence in the country has shown that only 32% of PLWHA on ART have suppressed viral loads (*UNAIDS, 2017*). Studies have indicated that lack of viral suppression is an indication of non-adherence to ART (*Byrd et al., 2019; Nasuuna et al., 2018; Robbins et al., 2014*). Therefore, understanding the level of adherence and factors related with non-adherence helps to prevent the future threat of widespread treatment resistant HIV strains. We, therefore, aimed at investigating the magnitude and associated factors with non-adherence to ART among adult people living with HIV and on treatment in the Benishangul-Gumuz Regional State (hereafter Benishangul-Gumuz) in northwest Ethiopia. Our results will inform implementation of strategic programs and practices to improve ART adherence to achieve optimal HIV care in the country.

METHODS

Study design and population

A cross-sectional study was conducted from mid-December 2016 to February 2017 in Benishangul-Gumuz, one of the nine regional states of the Federal Democratic Republic of Ethiopia located in the northwestern part of the country. The study was conducted among two referral hospitals and three health centers which provide comprehensive HIV care services. These health facilities were purposely targeted as they are utilized by the majority of people living with HIV. During this study period, there were a total of 2,721 PLWHA who were on ART treatment follow-up among the selected facilities. Sample size for this study was determined using the formula for the estimation of single proportion, $n = \frac{(z)^2 p(1-p)}{d^2}$, where proportion (p) of 63% taken from the previous study in Ethiopia (*Tiyou et al., 2012*) margin of error (d) = 5%, and 95% confidence limit (Z = 1.96). By adding 10% to cater for non-response rate, a total of 394 respondents were enrolled into the study. This was allotted to the study sites proportionally to the number of case load at each facility. Finally, a simple random sampling technique using a sampling frame developed from the registration book of the patients was used to enrol respondents daily at each study sites. The study entry criteria were: (a) PLWHA who had received at least

Table 1 Study sites and sample contribution.

Name of health facility	Accessible population (M)	Samples (frequency, %) $n = \sum[(\frac{M}{N}) * 394]$
Hospital A	1,079	156 (40%)
Hospital B	921	133 (34)
Health Centre X	275	40 (10)
Health Centre Y	251	37 (9%)
Health centre Z	195	28 (7%)
Total (N)	2,721	394 (100%)

Notes.

M, the number of peoples living with HIV and AIDS receiving ART at each health facilities; N, total number of PLWHA receiving ART at the study sites during the study period; n, number of samples proportionally drawn from each study site.

one-month antiretroviral therapy and on follow-up at the selected health facilities, and (b) resided in Benishangul Gumuz Regional State for at least two years, (c) signed consent to participate in the study. Exclusion criteria: (a) below 18 years of age, (b) critically ill, (c) psychiatric health problem as previously diagnosed and confirmed. [Table 1](#) presents the number of accessible populations or PLWHA receiving ART during the study period and sample contribution among the study sites.

Data and measures

A structured questionnaire was used for face-to-face interviews with the study population during times participants were waiting to see ART service providers and for pharmacy refills. The study tool was checked for clarity and consistency among the selected health facilities out of the study sites by a group of health providers over a two-week period. The result assisted in ensuring the quality and consistency of the instrument. Training was given to data collectors about the objectives of the study, and how they could approach the patients to obtain their confidentiality. The collected data kept in locked cabinets and password protected computers during data processing and analysis. Data extraction forms were reviewed daily for completeness. The questionnaire contained different variables such as sociodemographic and economic characteristics, clinical and anthropometric information.

Antiretroviral therapy adherence

Antiretroviral treatment adherence is the level to which a person is taking medicine as prescribed by a physician and as per medical recommendations in relation to timing, dosing and consistency, and correctly taking the drugs in terms of right doses and right times ([Chaiyachati et al., 2014](#)). In this study, patient self-assessments were collected using the visual analogue scale (VAS) to measure primary outcome and adherence. VAS method was used to assess the adherence mostly in resource-limited settings and the most effective and feasible method available at the time the study, and shown to predict virological response in clinical trials and clinical routines ([Finitis et al., 2016](#); [Chkhartishvili et al., 2014](#)). With the VAS approach, a series of scales with values ranging from 0 to 100% is used to assess adherence during the past 30 days ([Chesney et al., 2000](#)). The participants were asked to place an “X” inside the box above the point showing the best guess about how

much of their current antiretroviral medications were taken in the past 30 days. Adherence at a VAS score of $\geq 95\%$ over the past 30 days was defined as adherent and less than 95% as non-adherent.

Outcome measures

To investigate the extent and factors associated with non-adherence to ART among people living with HIV and on treatment, we used socioeconomic indicators such as education, income, asset possession, employment status, dietary diversity, nutritional status and household food insecurity. Sociodemographic characteristics such as age, gender and marital status; clinical characteristics such as CD4 count, duration on ART and history of opportunistic infections were taken into consideration during the interviews and chart review for data collection. Inclusion of independent variables were based on literature reviews conducted, data availability, and theoretical relevance (*Tabachnick & Fidell, 2007*). We chose these independent variables because of extensive literature highlighting the relationship between these covariates and ART adherence (*Fonsah et al., 2017; Eyassu, Mothiba & Mbambo-Kekana, 2016; Alagaw et al., 2013; Peltzer & Pengpid, 2013; Bangsberg, 2006*).

Food insecurity is defined as the economic and social condition of limited or uncertain access to adequate food (*US Department of Agriculture & Economic Research Service, 2019*). The access component of household food insecurity was measured using the standard household food insecurity access scale (HFIAS) (*Coates, Swindale & Bilinsky, 2007*). This is a nine-item questionnaire assessing household food insecurity in the domains of anxiety about household food access, insufficient quality of food and insufficient food intake in the past 30 days. In this study, HFIAS was calculated by summing the score for all nine items and ranges from 0 to a maximum of 27. A higher HFIAS score indicates poor access to food and greater food insecurity. Respondents were assigned to discrete categories of food insecurity severity: (1) food secure, (2) mildly food insecure, (3) moderately food insecure, and (4) severely food insecure, which we dichotomized into food insecure versus food secure. The alpha value for internal consistency of the scale was 0.91.

Malnutrition is the condition that occurs when the body does not get enough nutrients (*US National Library of Medicine, 2019*). In this study, malnutrition refers to undernutrition. The researchers used anthropometric measurement such as weight and height to measure malnutrition. The participants' weight was measured by means of the Seka weight scale calibrated to the nearest 0.1 kg after removing heavy clothes. The participants' height was measured using the Seka measuring rod calibrated to the nearest 0.1 cm. The participants took off their shoes, stood erect and looked straight in the horizontal plane to measure their height. The body mass index (BMI) was calculated as weight in kilograms divided by the square of height in meters (kg/m^2). Participants with a BMI of less than $18.5 \text{ kg}/\text{m}^2$ were considered malnourished.

Household dietary diversity is the economic ability of a household to access a variety of foods during the past seven days. Household dietary diversity was measured based on the dietary measurement method (*Kennedy, Ballard & Dop, 2011; Swindale & Bilinsky, 2006*). Twelve questions were used to assess dietary diversity. Participants were asked to report the

frequency of consumption of each of the following 12 food groups: (1) cereals; (2) roots and tubers; (3) pulses and legumes/nuts; (4) vegetables; (5) fruits; (6) meat and poultry; (7) eggs; (8) fish and seafood; (9) milk and milk products; (10) oils and fats; (11) sugar and sweets; and (12) miscellaneous. Irrespective of the frequency, participants received '1' point if they consumed at least once during the last seven days of the foods within each subgroup, and '0' points if they never consumed the food. The household dietary diversity score (HDDS) was constructed as the sum of some food groups consumed over the past week, ranging from 0 to 12. A high value indicated a diversified diet. The mean household dietary diversity score in the study subjects was categorized as adequate if the dietary diversity score is nine or above, and inadequate, when the diversity score is below nine.

Asset possession refers to a household's possession of assets elicited by asking participants a series of 13 questions about household assets and housing characteristics such as housing quality (floor, walls and roof material), source of drinking water, type of toilet facility, presence of electricity, type of cooking fuel, and ownership of modern household durable goods and livestock (e.g., bicycle, television, radio, motorcycle, telephone, refrigerator, mattress, bed and mobile phone). Following the method of Filmer and Pritchett (*Filmer & Pritchett, 2001*), principal components analysis was applied to define the asset wealth index among the PLWHA, and based on wealth index score participants were sorted into three quintiles of relative asset lower, middle and upper. A high value indicates more asset possession.

Data analyses

To assess factors associated with non-adherence to ART, binary adherence variables using data from VAS was created. Both descriptive and analytic statistics were used. Binary logistic regression analysis was used to find associations between the independent variables and the outcome variable. The Chi-square test of independence was used to find factors for not taking medication. All the variables showing a significant association in a binary analysis at $p < 0.25$ were entered to multivariable logistic regression to identify factors which have statistically significant association. In the final multivariate analysis, the test was two-sided and p -value < 0.05 was considered statistically significant. All statistical analyses were conducted in IBM SPSS Statistics for Windows, Version 24.0 (IBM Corporation, Armonk, NY, USA).

Ethical approval and consent to participate

Before the study begins, ethical clearance was obtained from the Health Studies Higher Degrees Committee of the College of Human Sciences at the University of South Africa (UNISA) (REC 012714-039 NHERC). Further, permission to conduct the study (ethical clearance) was sought after authorised cooperation letter was written from UNISA Ethiopia regional learning centre to Benishangul Gumuz Regional State Health Bureau (BG-RHB) and thus obtained the ethical clearance. A support letter to conduct the study among the selected health facilities in the region was further written to those health facilities by BGR-RHB. Institutional consent and permission to conduct an interview with PLWHA among

the selected health facilities were sought. Authorities at the study sites were assured that any information that the researcher came across during the conduct of the research was not be disclosed to any interest groups that could jeopardize the participants' welfare in society and the concerned institution. The researcher abided the guideline of the institutions, which has the right to terminate the study if the safety and confidentiality of participants might be compromised. All participants provided written informed consent to indicate their voluntary participation in the study before administering any instrument. The researchers ensured that participants understood their rights of voluntary participation, anonymity and confidentiality. ART follow-up records were retrieved by the health providers working in the ART units, and personal information of the participants was kept confidential.

RESULTS

Three hundred and ninety-four study participants were recruited for this study. Only four participants declined or terminated to participate into the study. The majority (80%) of the participants were aged between 25 and 44 years. The women participants constituted (66.4%) of the total participants. Nearly (50%) of the participants were married and an equal number of the participants were divorced, widowed or single. About (36.4%) of them never attended school, while only (4%) attended college/university. The majority of the participants (91%) lived in urban areas. [Table 2](#) shows the characteristics of the study participants.

In terms of socioeconomic characteristics, most of the participants were poor as measured by the wealth index, a proxy measure for asset possession. About (60%) of them were in the lower asset tertiles. More than a quarter of them were unemployed (31%). A large number of the participants (67.7%) lived on a poor mean monthly income of 1,260 Ethiopian Birr (equivalent to US\$45), below the World Bank poverty threshold of US\$ 1.90/day (*The World Bank Group, 2019*). The vast majority of the study participants (76%) were food insecure and (60%) had a BMI of less than 18.5 kg/m². The mean CD4 count was 559 (SD = 319.6) cells/mm³, with a range of 60 to 1,914 cell/mm³. More than a quarter of them (33.6%) suffered from frequent opportunistic infection in the last three months.

Based on a 30-day adherence measurement, (39.7%) of the study participants were found non-adherent to ART. Of the food insecure participants, more than half (51.4%) was non-adherent; while (47.2%) of malnourished PLWHA, those whose BMI is below 18.5 kg/m² was non-adherent to ART.

Factors associated with non-adherence to antiretroviral therapy

As presented in [Table 3](#), in the binary logistic regression analysis age below 25 years, marital status, urban residential area, never been to school, lack of employment, food insecurity, BMI below 18.5 kg/m², household dietary diversity, poor asset possession, CD4 cell count below 350 cell/mm³, and history of opportunistic infections were found associated with non-adherence to ART. But subsequent multivariate logistic regression analysis showed age below 25 years, lack of employment, food insecurity, malnutrition and opportunistic infections were among the factors associated with the factors in non-adherence to ART.

Table 2 Characteristics of study participants, $N = 390$.

Characteristics	Non-adherent to ART ($n = 155$)	Adherent to ART ($n = 235$)
<i>Sociodemographic characteristics</i>		
Age		
Less than 25 years	19 (12.0)	14 (6.0)
25–35 years	76 (49.0)	106 (45.1)
More than 35 years	60 (38.7)	115 (48.9)
Gender:		
Female	110 (71.0)	149 (63.4)
Male	45 (29.0)	86 (36.6)
Marital status:		
Divorced/Widowed/Single	67 (43.2)	128 (54.5)
Married	88 (56.8)	107 (45.5)
Religious:		
Christian	117 (75.5)	184 (78.3)
Muslim	38 (24.5)	51 (21.7)
Residence area:		
Urban	149 (96.1)	206 (87.7)
Rural	6 (3.9)	29 (12.3)
<i>Socioeconomic factors</i>		
Education level:		
Never been to school	57 (36.8)	85 (36.2)
Primary level	69 (44.5)	97 (41.3)
Secondary level	27 (17.4)	39 (16.6)
College/University level	2 (1.3)	14 (6.0)
Employment status:		
Unemployed	70 (45.2)	52 (22.1)
Employed	85 (54.8)	183 (77.9)
Monthly income (in Ethiopian Birr)		
<1750	119 (76.8)	174 (74)
≥1750	36 (23.2)	61 (26)
Food security status:		
Food insecure	152 (98.1)	144 (61.3)
Food secure	3 (1.9)	91 (38.7)
BMI score (in kg/m^2):		
<18.5	111 (71.6)	124 (52.8)
≥18.5	44 (28.4)	111 (47.2)
Household dietary diversity:		
Inadequate	121 (78.1)	139 (59.1)
Adequate	34 (21.9)	96 (40.9)

(continued on next page)

Table 2 (continued)

Characteristics	Non-adherent to ART (n = 155)	Adherent to ART (n = 235)
Household wealth index score		
Lower	76 (49.0)	158 (67.2)
Middle	35 (22.6)	41 (17.4)
Upper	44 (28.4)	36 (15.3)
Clinical features		
Duration of ART initiation:		
Less than 12 months	13 (8.4)	24 (10.2)
1–5 years	75 (48.4)	80 (34.0)
5–10 years	56 (36.1)	112 (47.7)
>10 years	11 (7.1)	19 (8.1)
CD4 count (in Cell/mm ³):		
<350	54 (34.8)	54 (23.0)
350–500	26 (16.8)	55 (23.4)
>501	75 (48.4)	126 (53.6)
History of opportunistic infections in the last three months:		
Yes	76 (49.0)	55 (23.4)
No	79 (51.0)	180 (76.6)

Notes.

n, frequency in number; %, percentage.

Age groups of less than 25 years were found to be four-fold non-adherent to ART in comparison with older age groups (AOR: 4.30, 95% CI [1.39–3.35]; $p = 0.011$). Urban residential area is also found contributing for non-adherence to ART (AOR: 2.78, CI [1.23–7.09], $p = 0.043$). The odds of non-adherence to ART among unemployed PLWHA were almost twice higher than their peers who were employed (AOR: 1.75, 95% CI [1.05–2.91]; $p = 0.032$). Food insecurity was the other socioeconomic factor strongly found impeding ART adherence (AOR: 2.67, 95% CI [7.59–8.97]; $p < 0.0001$), and malnutrition was among factors deterring PLWHA to attain the required $\geq 95\%$ ART adherence level (AOR: 1.55, 95% CI [1.94–2.56]; $p = 0.027$). Recurrent episodes of opportunistic infection (AOR: 1.81, 95% CI [1.11–2.97]; $p = 0.018$) was a clinical factor contributing to non-adherence to ART.

Reasons for missing medication

The prevalence of non-adherence to ART in this study was found to be (39.7%). Among these, (83%) of them missed twice a month while (4.4%) missed three times or more a month. Four reasons were reported by the study participants as important reasons for missing their medications: (i) demanding work or household responsibilities (33%), (ii) ran out of medication (34%), (iii) unable to take without food (22%), and (iv) forgetfulness (11%). Table 4 shows self-reported reasons for missing medications.

Table 3 Bivariable and multivariable analysis of factors associated with non-adherence to ART among peoples living with HIV and AIDS attending antiretroviral therapy in Benishangul Gumuz Region, Ethiopia, 2019.

Characteristics	Bivariable regression analysis			Multivariable regression analysis		
	COR	95% CI	p-value	AOR	95% CI	p-value
Sociodemographic characteristics						
Age						
Less than 25 years	2.6	[1.22, 5.55]	0.013	4.30	[1.39, 3.35]	0.011
25–35 years	1.4	[0.90, 2.10]	0.157	1.23	[0.75, 2.02]	0.411
More than 35 years	Ref					
Gender:						
Female	0.70	[0.46, 1.10]	0.120	1.09	[0.63, 1.09]	0.67
Male	Ref					
Marital status:						
Divorced/Widowed/Single	1.60	[1.04, 2.36]	0.030	0.63	[0.39, 1.01]	0.054
Married	Ref					
Religious:						
Christian	1.17	[0.71, 1.90]	0.520			
Muslim	Ref					
Residence area:						
Urban	3.50	[1.42, 8.60]	0.007	2.78	[1.23, 7.09]	0.043
Rural	Ref					
Socioeconomic factors						
Education level:						
Never been to school	4.70	[1.03, 21.0]	0.046	1.07	[0.17, 6.71]	0.38
Primary level	4.90	[1.1, 22.0]	0.038	1.44	[0.23, 0.89]	0.63
Secondary level	4.80	[1.02, 23]	0.047	1.42	[0.21, 9.46]	0.71
College/University level	Ref					
Employment status:						
Unemployed	2.89	[1.86, 4.50]	<0.001	1.75	[1.05, 2.91]	0.032
Employed	Ref					
Monthly income (in Ethiopian Birr)						
<1,750	0.86	[0.54, 1.39]	0.540			
≥1,750	Ref					
Food security status:						
Food insecure	3.2	[9.9, 11.3]	<0.0001	2.67	[7.59, 8.97]	<0.0001
Food secure	Ref					
BMI score (in kg/m ²):						
<18.5	2.3	[2.47, 4.48]	<0.0001	1.55	[1.94, 2.56]	0.027
≥18.5	Ref					
Household dietary diversity:						
Inadequate	2.5	[1.60, 3.90]	<0.0001	0.93	[0.51, 1.68]	0.81
Adequate	Ref					

(continued on next page)

Table 3 (continued)

Characteristics	Bivariable regression analysis			Multivariable regression analysis		
	COR	95% CI	p-value	AOR	95% CI	p-value
Household wealth index score						
Lower	1.8	[1.05, 3.01]	0.03	0.63	[0.33, 1.13]	0.12
Middle	2.5	[1.51, 4.27]	0.00	0.72	[0.35, 1.47]	0.36
Upper	Ref					
Clinical features						
Duration of ART initiation:						
Less than 12 months	0.94	[0.34, 2.50]	0.890			
1–5 years	1.60	[0.72, 3.60]	0.240			
5–10 years	0.86	[0.39, 1.90]	0.720			
>10 years	Ref					
CD4 count (in Cell/mm ³):						
<350	1.68	[1.10, 2.60]	0.032	1.1	[0.57, 1.78]	0.92
350–500	0.79	[0.46, 1.40]	0.410	0.77	[0.39, 1.46]	0.42
>501	Ref					
History of opportunistic infections in the last three months:						
Yes	3.20	[2.10, 4.90]	<0.001	1.81	[1.11, 2.97]	0.018
No	Ref					

Notes.

COR, crude odds ratio; AOR, adjusted odds ratio; OR, odds ratio; CI, confidence interval; Ref, reference.

Table 4 Self-reported reason for missing ART among peoples living with HIV and AIDS attending antiretroviral therapy in Benishangul Gumuz Region, Ethiopia, 2019.

Characteristics	Frequency	Percent (%)	χ^2	p-value
ART adherence:				
Adherent	235	60.3		
Non-adherent	135	39.7		
Reasons for not taking medication in the last seven days:				
Demanding work or household responsibilities	44	33.0		
Ran out of medication	46	34.0	3.01	0.009
Unable to take without food	30	22.0		
Forgetfulness	15	11.0		
Frequency of missing in the last seven days:				
Two times	112	83.0	5.48	0.043
Three times	17	12.6		
Above three	6	4.4		
VAS adherence score:				
M = 86.40				
SD = 14.94				

Notes.

χ^2 , Chi-square test; VAS, Visual Analogue Scale.

DISCUSSION

In the current study, the factors associated with non-adherence to ART among PLWHA in Benishangul-Gumuz were examined. Our findings endorse higher prevalence of ART non-adherence, with only 60.3% of our study participants remaining adherent to ART signposts the requirements of more efforts towards commitment to and realization of a 90-90-90 HIV treatment initiative. The study found age groups below 25 years were non-adherent to ART. Analogous to this finding, the results of a meta-analysis study (*Ghidei et al., 2013*) showed a higher risk of non-adherence among young PLWHA while older PLWHA had reduced risk of non-adherence to ART. Another study in Canada found a higher proportion of non-adherence among young PLWHA (*Hadland et al., 2012*). Youths are willing to engage in greater risky sexual behaviors and may also be more likely to assume the risks of poor HIV medication compliance, thus youth tailored behavioral change interventions that address the special emotional and developmental concerns of younger PLWHA could be designed.

Strong plausible evidence of the association between non-adherence to ART and food insecurity was found. Food insecure PLWHA were found twice more likely non-adherence to ART than their food secure counterparts. Studies in both developing and developed countries have culminated the strong association between non-adherence to ART and food insecurity effects. For example, studies in Lake Victoria, Kenya (*Nagata et al., 2012*) and in Atlanta, USA (*Pellowski et al., 2016*) showed that food insecure PLWHA miss doses of their daily medication. Similarly, studies in Jimma, southwest Ethiopia (*Tiyou et al., 2012*), and Congo (*Musumari et al., 2014*), and established a strong association between non-adherence to ART and food insecurity. Since non-adherence to ART can risk developing viral resistance (*Weiser et al., 2012; Musumari et al., 2014*) and food insecurity poses a significant challenge to the HIV/AIDS response (*Pellowski et al., 2016; Federal Ministry of Health of Ethiopia, 2015; Kalichman et al., 2014*).

In similar with the findings of other studies in Ethiopia (*Berhe, Tegabu & Alemayehu, 2013*), the current study found malnourished PLWHA were non-adherent to ART. This is also reported in another matched case-control study in the Central Tigray Zone, Ethiopia that revealed malnutrition was among the independent factors associated with lack of adherence to ART (*Weldehaweria et al., 2017*). Absence of nutritional support fuels the synergistic co-existence of HIV and malnutrition that has a perverse effect on ART adherence and the wellbeing of PLWHA (*Mekuria et al., 2015; Sicotte et al., 2014; Berhe, Tegabu & Alemayehu, 2013*). Therefore, nutritious food support is highly regarded as a viable option to increase weight gain (*Audain et al., 2015*) and improve treatment outcome (*Chaiyachati et al., 2014; Tiyou et al., 2012*).

Our findings are similar to the study by Nachega et al., who found a strong association between non-adherence to ART and lack of employment among PLWHA living both in low-income and in high-income countries (*Nagata et al., 2012*). In opposition to this finding, unemployed PLWHA in Nigeria had better ART adherence than employed people and rarely missed their ART medications (*Okoronkwo et al., 2013*). The possible explanation for adherence among unemployed PLWHA might be time freedom to take

their medications without forgetting being a concern. Also, they are more privileged to take their medications freely without any psychological impacts that put them under stress such as the workplace environment (*Okoronkwo et al., 2013*). Education status, dietary diversity and asset possession were among the alterable socioeconomic factors found influencing adherence to ART, thus developing effective adherence interventions to such alterable barriers are important in achieving an optimum adherence level.

Opportunistic infections were found to affect the treatment outcome of PLWHA. According to the results of this study, those who suffered recurrent illness within the last three months were twice more likely to miss their treatment and be non-adherent to ART. This finding is consistent with the findings of other studies that reported the presence of a significant association between treatment for HIV and other infections and adherence to ART (*Eyassu, Mothiba & Mbambo-Kekana, 2016*). Corresponding to this finding, non-adherence to ART as a result of opportunistic infection was reported among PLWHA in Yaounde, Cameroon (*Fonsah et al., 2017*) and the Republic of Korea (*Kim et al., 2018*). The reason for non-adherence to ART among sick PLWHA include the pill burden and poor absorption (*Fonsah et al., 2017; Eyassu, Mothiba & Mbambo-Kekana, 2016*).

Unlike the study of *Schafer et al. (2017)*, the findings of the current study reveals urban residential area as a risk factor for non-adherence to ART. This findings are consistent with the evidence that urban residences, as a result of dynamic socioeconomic alteration, may lack the desired adherence level (*Belayihun & Negus, 2015*). Similar to this finding, a study at Gonder University in northwest Ethiopia revealed that urban residential areas are associated with factors of non-adherence to ART (*Molla et al., 2018*). To alleviate the adherence issues both in urban and rural setups, a robust and sustainable response has to be foresighted, including using community volunteers to re-engage patients in care and helping them to remain in care, and empowering PLWHA in rural areas to regularly visit their healthcare provider (*Kaihin et al., 2015; Campbell et al., 2012*).

The current study has several limitations: first, even if an attempt was made to minimize social desirability bias, it might be subjected to recall and social desirability biases as ART adherence, food security and dietary diversity were assessed based on the self-reported measure. Second, the study design was cross-sectional in nature and hence cannot be used to determine causality. Malnutrition can be both a cause and effect of non-adherence to ART. Hence, future longitudinal studies are warranted to ascertain the causal directions of these variables.

CONCLUSIONS

The current study provides empirical findings on the factors associated with non-adherence to ART and the extent of non-adherence in Benishangul-Gumuz. The magnitude of non-adherence is as high as (39.7%) in the region. High prevalence of non-adherence is strongly associated with young age below 25 years, urban residential area, lack of employment, food insecurity, malnutrition and opportunistic infections. Therefore, improving ART adherence requires the adoption of innovative youth-centered adherence enhancing strategies, creation of employment opportunities, devising follow-up and mitigation of

recurrent opportunistic illness, and integrating nutrition and food security interventions with the HIV/AIDS program that can be implemented through the multisectoral approach to bridge the resource gap and improve the treatment outcome across all setups.

List of abbreviations

AIDS	Acquired Immune Deficiency Syndrome
ART	Antiretroviral Treatment
BG-RHB	Benishangul-Gumuz Regional State Health Bureau
BMI	Body Mass Index
HDDS	Household Dietary Diversity score
HFIAS	Household Food Insecurity Access Scale score
HIV	Human Immunodeficiency Virus
PLWHA	people living with HIV and AIDS
UNISA	University of South Africa
WFP	United Nations World Food Programme

ACKNOWLEDGEMENTS

We are grateful to all the research assistants, the data entry clerks and the PLWHA who voluntarily participated in the survey; and to the University of South Africa for the support provided to us.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The authors received no funding for this work.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Fikadu Tadesse Nigusso conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Azwihangwisi Helen Mavhandu-Mudzusi performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

Human Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

Health Studies Higher Degrees Committee of the College of Human Sciences at the University of South Africa approved the study (Ref: REC 012714-039 NHERC).

Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

Health Studies Higher Degrees Committee of the College of Human Sciences at the University of South Africa approved the study (Ref: REC 012714-039 NHERC).

Data Availability

The following information was supplied regarding data availability:

Raw data is available as a [Supplemental File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8558#supplemental-information>.

REFERENCES

- Abera A, Fenti B, Tesfaye T, Balcha F. 2015.** Factors influencing adherence to antiretroviral therapy among people living With HIV/AIDS at ART clinic in jimma university teaching hospital, Southwest Ethiopia. *Journal of Pharmacological Reports* 1:1–6 DOI 10.1007/s40495-015-0016-z.
- Alagaw A, Godana W, Taha M, Dejene T. 2013.** Factors associated with antiretroviral treatment adherence among adult patients in wolaita soddo hospital. *Journal of Tropical Diseases & Public Health* 1:1–7 DOI 10.4172/2329-891X.1000125.
- Audain KA, Zotor FB, Amuna P, Ellahi B. 2015.** Food supplementation among HIV-infected adults in Sub-Saharan Africa: impact on treatment adherence and weight gain. *The Proceedings of the Nutrition Society* 74:517–525 DOI 10.1017/S0029665115000063.
- Bangsberg DR. 2006.** Less than 95% adherence to nonnucleoside reverse-transcriptase inhibitor therapy can lead to viral suppression. *Clinical Infectious Diseases* 43:939–941 DOI 10.1086/507526.
- Basti BD, Mahesh V, Bant DD, Bathija GV. 2017.** Factors affecting antiretroviral treatment adherence among people living with human immunodeficiency virus/acquired immunodeficiency syndrome: a prospective study. *Journal of Family Medicine and Primary Care* 3:482–486 DOI 10.4103/2249-4863.222014.
- Belayihun B, Negus R. 2015.** Antiretroviral treatment adherence rate and associated factors among people living with HIV in Dubti hospital, Afar Regional State, East Ethiopia. *International Scholarly Research Notices* 2015:Article 187360 DOI 10.1155/2015/187360.
- Berhe N, Tegabu D, Alemayehu M. 2013.** Effect of nutritional factors on adherence to antiretroviral therapy among HIV-infected adults: a case control study in Northern Ethiopia. *BMC Infectious Diseases* 13:233 DOI 10.1186/1471-2334-13-233.
- Byrd KK, Hou JG, Hazen R, Kirkham H, Suzuki S, Clay PG, Bush T, Camp NM, Weidle PJ, Delpino A. 2019.** Antiretroviral adherence level necessary for HIV viral

- suppression using real-world data. *Journal of Acquired Immune Deficiency Syndromes* 82(3):245–251 DOI 10.1097/QAI.0000000000002142.
- Campbell C, Skovdal M, Mupambireyi Z, Madanhire C, Nyamukapa C, Gregson S. 2012.** Building adherence-competent communities: factors promoting children’s adherence to anti-retroviral HIV/AIDS treatment in rural Zimbabwe. *Health & Place* 18:123–131 DOI 10.1016/j.healthplace.2011.07.008.
- Chaiyachati KH, Ogbuoji O, Price M, Suthar AB, Negussie EK, Bärnighausen T. 2014.** Interventions to improve adherence to antiretroviral therapy: a rapid systematic review. *AIDS* 28(Suppl 2):S187–S204 DOI 10.1097/QAD.0000000000000252.
- Chauhan CK, Lakshmi PVM, Sagar V, Sharma A, Arora SK, Kumar R. 2019.** Primary HIV drug resistance among recently infected cases of HIV in North-West India. *AIDS Research and Treatment* 2019:1–8 DOI 10.1155/2019/1525646.
- Chesney MA, Ickovics JR, Chambers DB, Gifford AL, Neidig J, Zwickl B, Wu AW. 2000.** Self-reported adherence to antiretroviral medications among participants in HIV clinical trials: The AACTG Adherence Instruments. *AIDS Care* 12:255–266 DOI 10.1080/09540120050042891.
- Chkhartishvili N, Rukhadze N, Svanidze M, Sharvadze L, Dehovitz JA, Tsertsvadze T, McNutt L-A, Del Rio C. 2014.** Evaluation of multiple measures of antiretroviral adherence in the Eastern European country of Georgia. *Journal of the International AIDS Society* 17(1):18885 DOI 10.7448/IAS.17.1.18885.
- Coates J, Swindale A, Bilinsky . 2007.** Household Food Insecurity Access Scale (HFAS) for measurement of food access: indicator guide: version 3: (576842013-001) DOI 10.1037/e576842013-001.
- Dalhatu I, Onotu D, Odafe S, Abiri O, Debem H, Agolory S, Shiraishi RW, Auld AF, Swaminathan M, Dokubo K, Ngige E, Asadu C, Abatta E, Ellerbrock TV. 2016.** Outcomes of Nigeria’s HIV/AIDS treatment program for patients initiated on antiretroviral treatment between 2004–2012. *PLOS ONE* 11:e0165528 DOI 10.1371/journal.pone.0165528.
- Deribew A, Biadgilign S, Deribe K, Dejene T, Tessema GA, Melaku YA, Lakew Y, Amare AT, Bekele T, Abera SF, Kumsa A, Assefa Y, Glenn SD, Carter A, Misganaw A, Wang H. 2019.** The burden of HIV/AIDS in Ethiopia from 1990 to 2016: evidence from the global burden of diseases 2016 study, vol. 29. 10.
- Eyassu MA, Mothiba TM, Mbambo-Kekana NP. 2016.** Adherence to antiretroviral therapy among HIV and AIDS patients at the Kwa-Thema clinic in Gauteng Province, South Africa. *African Journal of Primary Health Care & Family Medicine* 8(2):924 DOI 10.4102/phcfm.v8i2.924.
- Federal Ministry of Health of Ethiopia. 2015.** *Health sector transformation plan 2015/16-2019/20 (2008-2012 EFY)*. Addis Ababa, Ethiopia: Ministry of Health.
- Filmer D, Pritchett LH. 2001.** Estimating wealth effects without expenditure data—or tears: an application to educational enrollments in states of India. *Demography* 38:115–132.
- Finitsis DJ, Pellowski JA, Huedo-Medina TB, Fox MC, Kalichman SC. 2016.** Visual analogue scale (VAS) measurement of antiretroviral adherence in people living

- with HIV (PLWH): a meta-analysis. *Journal of Behavioral Medicine* **39**:1043–1055
DOI [10.1007/s10865-016-9770-6](https://doi.org/10.1007/s10865-016-9770-6).
- Fonsah JY, Njamnshi AK, Kouanfack C, Qiu F, Njamnshi DM, Tagny CT, Nchindap E, Kenmogne L, Mbanya D, Heaton R, Kanmogne GD. 2017.** Adherence to Antiretroviral Therapy (ART) in Yaoundé-Cameroon: association with opportunistic infections, depression, ART regimen and side effects. *PLOS ONE* **12**:e0170893
DOI [10.1371/journal.pone.0170893](https://doi.org/10.1371/journal.pone.0170893).
- Ghidei L, Simone M, Salow M, Zimmerman K, Paquin AM, Skarf LM, Kostas T, Rudolph JL. 2013.** Aging, antiretrovirals, and adherence: a meta analysis of adherence among older HIV-infected individuals. *Drugs & Aging* **30**:809–819
DOI [10.1007/s40266-013-0107-7](https://doi.org/10.1007/s40266-013-0107-7).
- Granich R, Kahn JG, Bennett R, Holmes CB, Garg N, Serenata C, Sabin ML, Makhoulf-Obermeyer C, Mack CDeFilippo, Williams P, Jones L, Smyth C, Kutch KA, Ying-Ru L, Vitoria M, Souteyrand Y, Crowley S, Korenromp EL, Williams BG. 2012.** Expanding ART for treatment and prevention of HIV in South Africa: estimated cost and cost-effectiveness 2011-2050. *PLOS ONE* **7**:e30216
DOI [10.1371/journal.pone.0030216](https://doi.org/10.1371/journal.pone.0030216).
- Hadland SE, Milloy M-J, Kerr T, Zhang R, Guillemi S, Hogg RS, Montaner JS, Wood E. 2012.** Young age predicts poor antiretroviral adherence and viral load suppression among injection drug users. *AIDS Patient Care and STDs* **26**:274–280
DOI [10.1089/apc.2011.0196](https://doi.org/10.1089/apc.2011.0196).
- Hansana V, Sanchaisuriya P, Durham J, Sychareun V, Chaleunvong K, Boonyaleepun S, Schelp FP. 2013.** Adherence to Antiretroviral Therapy (ART) among people living with HIV (PLHIV): a cross-sectional survey to measure in Lao PDR. *BMC Public Health* **13**:617 DOI [10.1186/1471-2458-13-617](https://doi.org/10.1186/1471-2458-13-617).
- Iacob SA, Iacob DG, Jugulete G. 2017.** Improving the adherence to antiretroviral therapy, a difficult but essential task for a successful hiv treatment—clinical points of view and practical considerations. *Frontiers in Pharmacology* **8**:1–12
DOI [10.3389/fphar.2017.00831](https://doi.org/10.3389/fphar.2017.00831).
- Kaihin R, Kasatpibal N, Chitreechuer J, Grimes RM. 2015.** Effect of an empowerment intervention on antiretroviral drug adherence in Thai youth. *Behavioral Medicine* **41**:186–194 DOI [10.1080/08964289.2014.911717](https://doi.org/10.1080/08964289.2014.911717).
- Kalichman SC, Grebler T, Amaral CM, McKerney M, White D, Kalichman MO, Cherry C, Eaton L. 2014.** Food insecurity and antiretroviral adherence among HIV positive adults who drink alcohol. *Journal of Behavioral Medicine* **37**:1009–1018
DOI [10.1007/s10865-013-9536-3](https://doi.org/10.1007/s10865-013-9536-3).
- Karade S, Chaturbhuj DN, Sen S, Joshi RK, Kulkarni SS, Shankar S, Gangakhedkar RR. 2018.** HIV drug resistance following a decade of the free antiretroviral therapy programme in India: a review. *International Journal of Infectious Diseases* **66**:33–41
DOI [10.1016/j.ijid.2017.10.020](https://doi.org/10.1016/j.ijid.2017.10.020).
- Kennedy G, Ballard T, Dop M-C. 2011.** *Guidelines for measuring household and individual dietary diversity*. Rome: FAO.

- Kim J, Lee E, Park B-J, Bang JH, Lee JY. 2018.** Adherence to antiretroviral therapy and factors affecting low medication adherence among incident HIV-infected individuals during 2009–2016: a nationwide study. *Scientific Reports* **8**:3133 DOI [10.1038/s41598-018-21081-x](https://doi.org/10.1038/s41598-018-21081-x).
- Liping M, Peng X, Haijiang L, Lahong J, Fan LH, Yuhua R. 2015.** Quality of life of people living with HIV/AIDS: a cross-sectional study in Zhejiang Province, China. *PLOS ONE* **10**(8):e0135705 DOI [10.1371/journal.pone.0135705](https://doi.org/10.1371/journal.pone.0135705).
- Mekuria LA, Sprangers MAG, Prins JM, Yalew AW, Nieuwkerk PT. 2015.** Health-related quality of life of HIV-infected adults receiving combination antiretroviral therapy in Addis Ababa. *AIDS Care* **27**:934–945 DOI [10.1080/09540121.2015.1020748](https://doi.org/10.1080/09540121.2015.1020748).
- Mohammed AY, Ahmed MB, Tefera TB. 2015.** Assessment of factors affecting art adherence among people living with human immune virus in bale robe hospital, South East Ethiopia. *American Journal of Public Health Research* **3**:60–67 DOI [10.12691/ajphr-3-2-5](https://doi.org/10.12691/ajphr-3-2-5).
- Molla AA, Gelagay AA, Mekonnen HS, Teshome DF. 2018.** Adherence to antiretroviral therapy and associated factors among HIV positive adults attending care and treatment in University of Gondar Referral Hospital, Northwest Ethiopia. *BMC Infectious Diseases* **18**:Article 266 DOI [10.1186/s12879-018-3176-8](https://doi.org/10.1186/s12879-018-3176-8).
- Musumari PM, Wouters E, Kayembe PK, Kiumbu Nzita M, Mbikayi SM, Suguimoto SP, Techasrivichien T, Lukhele BW, El-saaidi C, Piot P, Ono-Kihara M, Kihara M. 2014.** Food insecurity is associated with increased risk of non-adherence to antiretroviral therapy among hiv-infected adults in the democratic republic of congo: a cross-sectional study. *PLOS ONE* **9**:e85327 DOI [10.1371/journal.pone.0085327](https://doi.org/10.1371/journal.pone.0085327).
- Nachea J, Marconi VC, Van Zyl GU, Gardner EM, Preiser W, Hong SY, Mills EJ, Gross R. 2011.** HIV treatment adherence, drug resistance, virologic failure: evolving concepts. *Infectious Disorders - Drug Targets* **11**:167–174 DOI [10.2174/187152611795589663](https://doi.org/10.2174/187152611795589663).
- Nagata JM, Magerenge RO, Young SL, Oguta JO, Weiser SD, Cohen CR. 2012.** Social determinants, lived experiences, and consequences of household food insecurity among persons living with HIV/AIDS on the shore of Lake Victoria, Kenya. *AIDS Care* **24**:728–736 DOI [10.1080/09540121.2011.630358](https://doi.org/10.1080/09540121.2011.630358).
- Nasuuna E, Kigozi J, Babirye L, Muganzi A, Sewankambo NK, Nakanjako D. 2018.** Low HIV viral suppression rates following the intensive adherence counseling (IAC) program for children and adolescents with viral failure in public health facilities in Uganda. *BMC Public Health* **18**:1048 DOI [10.1186/s12889-018-5964-x](https://doi.org/10.1186/s12889-018-5964-x).
- Oguntibeju O. 2012.** Quality of life of people living with HIV and AIDS and antiretroviral therapy. *HIV/AIDS - Research and Palliative Care* **2012**:17–124 DOI [10.2147/HIV.S32321](https://doi.org/10.2147/HIV.S32321).
- Okoronkwo I, Okeke U, Chinweuba A, Iheanacho . 2013.** Nonadherence factors and sociodemographic characteristics of hiv-infected adults receiving antiretroviral therapy in nnamdi Azikiwe University Teaching Hospital, Nnewi, Nigeria. Available at <https://www.hindawi.com/journals/isrn/2013/843794/> (accessed on 8 April 2019).

- Paterson DL, Swindells S, Mohr J, Brester M, Vergis EN, Squier C, Wagener MM, Singh N. 2000.** Adherence to protease inhibitor therapy and outcomes in patients with HIV infection. *Annals of Internal Medicine* **133**:21–30
DOI [10.7326/0003-4819-133-1-200007040-00004](https://doi.org/10.7326/0003-4819-133-1-200007040-00004).
- Pellowski JA, Kalichman SC, Cherry S, Conway-Washington C, Cherry C, Grebler T, Krug L. 2016.** The daily relationship between aspects of food insecurity and medication adherence among people living with hiv with recent experiences of hunger. *Annals of Behavioral Medicine: A Publication of the Society of Behavioral Medicine* **50**:844–853 DOI [10.1007/s12160-016-9812-x](https://doi.org/10.1007/s12160-016-9812-x).
- Peltzer K, Pengpid S. 2013.** Socioeconomic factors in adherence to HIV therapy in low- and middle-income countries. *Journal of Health, Population, and Nutrition* **31**:150–170.
- Robbins RN, Spector AY, Mellins CA, Remien RH. 2014.** Optimizing ART adherence: update for HIV treatment and prevention. *Current HIV/AIDS reports* **11**:423–433
DOI [10.1007/s11904-014-0229-5](https://doi.org/10.1007/s11904-014-0229-5).
- Samuel Edward K, Maseke Richard M, Basiliana E, Geoffrey Nimrod S, Debora Charles K. 2018.** Factors influencing adherence to antiretroviral therapy among HIV infected patients in Nyamagana-Mwanza, Northern Tanzania: a cross sectional study. *International Archives of Medical Microbiology* **1**:002 DOI [10.23937/iamm-2017/1710002](https://doi.org/10.23937/iamm-2017/1710002).
- Schafer KR, Albrecht H, Dillingham R, Hogg RS, Jaworsky D, Kasper K, Loutfy M, MacKenzie LJ, McManus KA, Oursler KAK, Rhodes SD, Samji H, Skinner S, Sun CJ, Weissman S, Ohl ME. 2017.** The continuum of HIV care in rural communities in the United States and Canada: what is known and future research directions. *Journal of Acquired Immune Deficiency Syndromes* **75**:35–44
DOI [10.1097/QAI.0000000000001329](https://doi.org/10.1097/QAI.0000000000001329).
- Sicotte M, Langlois ÉV, Aho J, Ziegler D, Zunzunegui MV. 2014.** Association between nutritional status and the immune response in HIV, + patients under HAART: protocol for a systematic review. *Systematic Reviews* **3**:9 DOI [10.1186/2046-4053-3-9](https://doi.org/10.1186/2046-4053-3-9).
- Swindale A, Bilinsky P. 2006.** Household Dietary Diversity Score (HDDS) for measurement of household food access: indicator guide (Version 2). Available at https://www.fantaproject.org/sites/default/files/resources/HDDS_v2_Sep06_0.pdf (accessed on 12 June 2019).
- Tabachnick B, Fidell L. 2007.** *Using multivariate statistics*. 5th edition. New York: Allyn and Bacon.
- The Federal Democratic Republic of Ethiopia Ministry of Health. 2015.** Health Sector Transformation Plan (HSTP) 2015/16–2019/20 (2008–2015). In: *The Federal Democratic Republic of Ethiopia Ministry of Health*.
- The World Bank Group. 2019.** Poverty overview: Ethiopia. Available at <https://www.worldbank.org/en/topic/poverty/overview> (accessed on 8 April 2019).
- Tiyou A, Belachew T, Alemseged F, Biadgilign S. 2012.** Food insecurity and associated factors among HIV-infected individuals receiving highly active antiretroviral therapy in Jimma zone Southwest Ethiopia. *Nutrition Journal* **11**:Article 51
DOI [10.1186/1475-2891-11-51](https://doi.org/10.1186/1475-2891-11-51).

- UNAIDS. 2011. *A new investment framework for the global HIV response*. Geneva: UNAIDS.
- UNAIDS. 2015. How AIDS changed everything —MDG6: 15 years, 15 lessons of hope from the AIDS response. Available at http://www.unaids.org/en/resources/documents/2015/MDG6_15years-15lessonsfromtheAIDSresponse (accessed on 6 April 2019).
- UNAIDS. 2017. UNAIDS DATA 2017. Geneva: UNAIDS.
- UNAIDS. 2018. Global HIV & AIDS statistics —2018 fact sheet. Available at <https://www.unaids.org/en/resources/fact-sheet> (accessed on 17 April 2019).
- US Department of Agriculture, Economic Research Service. 2019. Definitions of food security. Available at <https://www.ers.usda.gov/topics/food-nutrition-assistance/food-security-in-the-us/definitions-of-food-security.aspx> (accessed on 8 April 2019).
- US National Library of Medicine. 2019. Malnutrition: Definition. Available at <https://medlineplus.gov/ency/article/000404.htm> (accessed on 8 April 2019).
- Wang EA, McGinnis KA, Fiellin DA, Goulet JL, Bryant K, Gibert CL, Leaf DA, Mattocks K, Sullivan LE, Vogenthaler N, Justice AC. 2011. Food insecurity is associated with poor virologic response among HIV-infected patients receiving antiretroviral medications. *Journal of General Internal Medicine* 26:1012–1018 DOI 10.1007/s11606-011-1723-8.
- Wasti SP, Simkhada P, Randall J, Freeman JV, Teijlingen Evan. 2012. Factors influencing adherence to antiretroviral treatment in nepal: a mixed-methods study. *PLOS ONE* 7:e35547 DOI 10.1371/journal.pone.0035547.
- Weiser SD, Tsai AC, Gupta R, Frongillo EA, Kawuma A, Senkungu J, Hunt PW, Emenyonu NI, Mattson JE, Martin JN, Bangsberg DR. 2012. Food insecurity is associated with morbidity and patterns of healthcare utilization among HIV-infected individuals in a resource-poor setting. *AIDS* 26:67–75 DOI 10.1097/QAD.0b013e32834cad37.
- Weldehaweria NB, Abreha EH, Weldu MG, Misgina KH. 2017. Psychosocial correlates of nutritional status among people living with HIV on antiretroviral therapy: a matched case-control study in Central zone of Tigray, Northern Ethiopia. *PLOS ONE* 12:e0174082 DOI 10.1371/journal.pone.0174082.
- Williams BG, Lima V, Gouws E. 2011. Modelling the impact of antiretroviral therapy on the epidemic of HIV. *Current HIV Research* 9:367–382 DOI 10.2174/157016211798038533.
- World Health Organization. 2010. Antiretroviral drugs for treating pregnant women and preventing HIV infection in infants: recommendations for a public health approach. Geneva: World Health Organization.
- Young S, Wheeler A, McCoy S, Weiser SD. 2014. A review of the role of food insecurity in adherence to care and treatment among adult and pediatric populations living with HIV and AIDS. *AIDS and Behavior* 18:505–515 DOI 10.1007/s10461-013-0547-4.